arXiv:1410.2633v1 [math.NA] 9 Oct 2014

A new class of optimal four-point methods with convergence order 16 for solving nonlinear equations

Somayeh Sharifi[a*] Mehdi Salimi[b†] Stefan Siegmund[b‡] Taher Lotfi[c§]

[a]Young Researchers and Elite Club, Hamedan Branch, Islamic Azad University, Hamedan, Iran
[b]Center for Dynamics, Department of Mathematics, Technische Universität Dresden, 01062 Dresden, Germany
[c]Department of Mathematics, Hamedan Branch, Islamic Azad University, Hamedan, Iran

Abstract

We introduce a new class of optimal iterative methods without memory for approximating a simple root of a given nonlinear equation. The proposed class uses four function evaluations and one first derivative evaluation per iteration and it is therefore optimal in the sense of Kung and Traub's conjecture. We present the construction, convergence analysis and numerical implementations, as well as comparisons of accuracy and basins of attraction between our method and existing optimal methods for several test problems.

Keywords: Simple root, four-step iterative method, Kung and Traub conjecture, optimal order of convergence, computational efficiency.

1 Introduction

Solving nonlinear equations is a basic and extremely valuable tool in all fields in the sciences and in engineering. One can distinguish between two general approaches for solving nonlinear equations numerically, namely, one-step and multi-step methods. Multi-step methods overcome some computational issues encountered with one-step iterative methods, typically they allow us to achieve a greater accuracy with the same number of function evaluations. In this context an unproved conjecture by Kung and Traub [18] plays a central role, it states that an optimal multi-step method without memory which uses $n+1$ evaluations could achieve a convergence order of 2^n. Considering this conjecture, many optimal two-step and three-step methods have been presented. However, because of complexity in construction and development, optimal four-point methods are rare and can be considered as an active research problem.

*s.sharifi@iauh.ac.ir
†Corresponding author: mehdi.salimi@tu-dresden.de
‡stefan.siegmund@tu-dresden.de
§lotfi@iauh.ac.ir

Prominent optimal two-point methods have been introduced by Jarratt [15], King [17] and Ostrowski [26]. Some optimal three-point methods have been proposed by Chun and Lee [6], Cordero et al. [7]-[10], Khattri and Steihaug [16], Lotfi et al. [19]-[21], Petkovic et al. [27, 28] and Sharma et al. [30]. Neta [22] has presented methods with convergence orders 8 and 16. Babajee and Thukral [5] developed a four-point method with convergence order 16 based on the King family of methods. In [11]-[13] Geum and Kim provided three methods with convergence order 16 by using weight function methods.

We construct a new optimal class of four-point methods without memory which uses five function evaluations per iteration. The paper is organized as follows: Section 2 is devoted to the construction of the new optimal class. We first construct classes of optimal two-point and three-point methods and then utilize them in the three first steps of our new method. The section also includes convergence analysis of all these methods. Numerical performance and comparisons with other methods are illustrated in Section 3. A conclusion is provided in Section 4.

2 Main results: Construction, error and convergence analysis

This section deals with construction, and error and convergence analysis of our method. First, we try to introduce an optimal two-point class (this class contains no originality, and we review it here only for easy-reference in constructing the next two classes), then an optimal three-point class is presented, and, finally, our optimal four-point class will be developed.

2.1 Construction of an optimal two-point class

In this section we construct a new optimal two-point method for solving nonlinear equations which employs Newton's one-point method [26, 33] and suitable weight functions for evaluations at two points.

Newton's method [26, 33]

$$y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \quad (n \in \mathbb{N}_0 = \{0, 1, \dots\}), \tag{2.1}$$

where x_0 denotes the initial approximation of x^*, is of convergence order two. To increase the order of convergence, we add one Newton step to the method (2.1) to get

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ x_{n+1} = y_n - \dfrac{f(y_n)}{f'(y_n)}. \end{cases} \tag{2.2}$$

The method (2.2) uses four function evaluations to achieve order four, therefore the method is not an optimal two-point method. We modify (2.2) by approximating $f'(y_n)$ by

$$f'(y_n) \approx \frac{f'(x_n)}{G(t_n)},$$

using only the values $f(x_n)$, $f(y_n)$, and $f'(x_n)$. More precisely, we use the abbreviations $t_n = \frac{f(y_n)}{f(x_n)}$ and utilize Mathematica [14] to carefully choose the weight function $G : \mathbb{R} \to \mathbb{R}$ from a class of admissible functions such that the scheme

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ x_{n+1} = y_n - G(t_n) \cdot \dfrac{f(y_n)}{f'(x_n)}, \end{cases} \tag{2.3}$$

is of order four.

Theorem 1. *Assume that $f : D \subset \mathbb{R} \to \mathbb{R}$ is four times continuously differentiable and has a simple zero $x^* \in D$ and $G \in C^3\mathbb{R}$ is sufficiently continuously differentiable. If*

$$G_0 = 1, \quad G_1 = 2, \quad |G_2| < \infty,$$

where $G_i = \frac{d^i G(t_n)}{dt_n^i}|_0$ for $i = 0, 1, \ldots$, and the initial point x_0 is sufficiently close to x^*, then the sequence $\{x_n\}$ defined by (2.3) converges to x^* and the order of convergence is four.

Proof. Let $e_n := x_n - x^*$, $e_{n,y} := y_n - x^*$, and $c_n := \frac{f^{(n)}(x^*)}{n! f'(x^*)}$ for $n \in \mathbb{N}$. Using the fact that $f(x^*) = 0$, the Taylor expansion of f at x^* yields

$$f(x_n) = f'(x^*)(e_n + c_2 e_n^2 + c_3 e_n^3 + c_4 e_n^4) + O(e_n^5), \tag{2.4}$$

and expanding f' at x^* we get

$$f'(x_n) = f'(x^*)(1 + 2c_2 e_n + 3c_3 e_n^2 + 4c_4 e_n^3 + 5c_5 e_n^4) + O(e_n^5). \tag{2.5}$$

Therefore

$$\frac{f(x_n)}{f'(x_n)} = e_n - c_2 e_n^2 + 2(c_2^2 - c_3)e_n^3 + (-4c_2^3 + 7c_2 c_3 - 3c_4)e_n^4 + O(e_n^5),$$

and hence

$$e_{n,y} = y_n - x^* = c_2 e_n^2 + O(e_n^3).$$

For $f(y_n)$, we also have

$$f(y_n) = f'(x^*)\left(e_{n,y} + c_2 e_{n,y}^2 + c_3 e_{n,y}^3 + c_4 e_{n,y}^4\right) + O(e_{n,y}^5). \tag{2.6}$$

By (2.4) and (2.6), we obtain

$$t_n = \frac{f(y_n)}{f(x_n)} = c_2 e_n + (-3c_2^2 + 2c_3)e_n^2 + (8c_2^3 - 10c_2 c_3 + 3c_4)e_n^3 + O(e_n^4), \tag{2.7}$$

and expanding G at 0 yields

$$G(t_n) = G_0 + G_1 t_n + \frac{1}{2} G_2 t_n^2 + O(t_n^3). \tag{2.8}$$

By substituting (2.4)-(2.8) into (2.3), we obtain

$$e_{n+1} = x_{n+1} - x^* = R_2 e_n^2 + R_3 e_n^3 + R_4 e_n^4 + O(e_n^5),$$

where

$$\begin{aligned} R_2 &= -c_2(-1 + G_0), \\ R_3 &= -c_2^2(-2 + G_1), \\ R_4 &= -c_2 c_3 + c_2^3(5 - \frac{1}{2}G_2). \end{aligned}$$

In general $R_4 \neq 0$, however, by setting $R_2 = R_3 = 0$, the convergence order becomes four. Sufficient conditions are given by the following set of equations

$$\begin{aligned} G_0 = 1 \quad &\Rightarrow \quad R_2 = 0, \\ G_1 = 2 \quad &\Rightarrow \quad R_3 = 0, \\ |G_2| < \infty \quad &\Rightarrow \quad R_4 \neq 0, \end{aligned}$$

and the error equation becomes

$$e_{n+1} = R_4 e_n^4 + O(e_n^5),$$

which finishes the proof. □

2.2 Construction of an optimal three-point class

To increase the order of convergence, we add one Newton step to the method (2.3) to get

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - G(t_n) \cdot \dfrac{f(y_n)}{f'(x_n)}, \\ x_{n+1} = z_n - \dfrac{f(z_n)}{f'(z_n)}. \end{cases} \tag{2.9}$$

The method (2.9) uses five function evaluations and therefore the method is not an optimal three-point method. We modify (2.9) by approximating $f'(z_n)$ by

$$f'(z_n) \approx \frac{f'(x_n)}{H(t_n, s_n, u_n)},$$

using only the values $f(x_n)$, $f(y_n)$, $f(z_n)$ and $f'(x_n)$. More precisely, we use the abbreviations $t_n = \frac{f(y_n)}{f(x_n)}$, $s_n = \frac{f(z_n)}{f(y_n)}$, $u_n = \frac{f(z_n)}{f(x_n)}$ and utilize Mathematica [14] to carefully choose the weight functions $H : \mathbb{R}^3 \to \mathbb{R}$ from a class of admissible functions such that the scheme

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - G(t_n) \cdot \dfrac{f(y_n)}{f'(x_n)}, \\ x_{n+1} = z_n - H(t_n, s_n, u_n) \cdot \dfrac{f(z_n)}{f'(x_n)}, \end{cases} \tag{2.10}$$

is of order eight.

Theorem 2. *Assume that $f : D \subset \mathbb{R} \to \mathbb{R}$ is eight times continuously differentiable and has a simple zero $x^* \in D$ and $G \in C^3\mathbb{R}$ and $H : \mathbb{R}^3 \to \mathbb{R}$ are sufficiently often differentiable functions. If*

$$\begin{aligned} &G_2 = 10, \quad G_3 = -36, \\ &H_{0,0,0} = 1, \quad H_{1,0,0} = 2, \quad H_{0,1,0} = 1, \\ &H_{2,0,0} = 12, \quad H_{0,0,1} = 4, \quad H_{1,1,0} = 0, \end{aligned}$$

where $H_{i,j,k} = \frac{\partial^{i+j+k} H(t_n, s_n, u_n)}{\partial t_n^i \partial s_n^j \partial u_n^k}|_{(0,0,0)}$ for $i, j, k = 0, 1, 2, 3, \ldots$, and the initial point x_0 is sufficiently close to x^*, then the sequence $\{x_n\}$ defined by (2.10) converges to x^* and the order of convergence is eight.

Proof. Let $e_n := x_n - x^*$, $e_{n,y} := y_n - x^*$, $e_{n,z} := z_n - x^*$ and $c_n := \frac{f^{(n)}(x^*)}{n! f'(x^*)}$ for $n \in \mathbb{N}$. Using the fact that $f(x^*) = 0$, the Taylor expansion of f at x^* yields

$$f(x_n) = f'(x^*)(e_n + c_2 e_n^2 + c_3 e_n^3 + \ldots + c_8 e_n^8) + O(e_n^9), \tag{2.11}$$

and expanding f' at x^* we get

$$f'(x_n) = f'(x^*)(1 + 2c_2 e_n + 3c_3 e_n^2 + \ldots + 9c_9 e_n^8) + O(e_n^9). \tag{2.12}$$

Therefore

$$\begin{aligned} \frac{f(x_n)}{f'(x_n)} = {} & e_n - c_2 e_n^2 + 2(c_2^2 - c_3)e_n^3 + (-4c_2^3 + 7c_2c_3 - 3c_4)e_n^4 + (8c_2^4 - 20c_2^2c_3 + 6c_3^2 \\ & + 10c_2c_4 - 4c_5)e_n^5 + (-16c_2^5 + 52c_2^3c_3 - 28c_2^2c_4 + 17c_3c_4 + c_2(-33c_3^2 + 13c_5))e_n^6 \\ & + 2(16c_2^6 - 64c_2^4c_3 - 9c_3^3 + 36c_2^3c_4 - 46c_2c_3c_4 + 6c_4^2 + 9c_2^2(7c_3^2 - 2c_5) + 11c_3c_5)e_n^7 \\ & + (-64c_2^7 + 304c_2^5c_3 - 176c_2^4c_4 + 348c_2^2c_3c_4 + c_4(-75c_3^2 + 31c_5) \\ & + c_2^3(-408c_3^2 + 92c_5) + c_2(135c_3^3 - 64c_4^2 - 118c_3c_5))e_n^8 + O(e_n^9) \end{aligned}$$

and hence

$$\begin{aligned} e_{n,y} = y_n - x^* = {} & c_2 e_n^2 - 2(c_2^2 - c_3)e_n^3 + (4c_2^3 - 7c_2c_3 + 3c_4)e_n^4 - (8c_2^4 - 20c_2^2c_3 + 6c_3^2 + 10c_2c_4 - 4c_5)e_n^5 \\ & - (-16c_2^5 + 52c_2^3c_3 - 28c_2^2c_4 + 17c_3c_4 + c_2(-33c_3^2 + 13c_5))e_n^6 \\ & - 2(16c_2^6 - 64c_2^4c_3 - 9c_3^3 + 36c_2^3c_4 - 46c_2c_3c_4 + 6c_4^2 + 9c_2^2(7c_3^2 - 2c_5) + 11c_3c_5)e_n^7 \\ & - (-64c_2^7 + 304c_2^5c_3 - 176c_2^4c_4 + 348c_2^2c_3c_4 + c_4(-75c_3^2 + 31c_5) \\ & + c_2^3(-408c_3^2 + 92c_5) + c_2(135c_3^3 - 64c_4^2 - 118c_3c_5))e_n^8 + O(e_n^9) \end{aligned}$$

For $f(y_n)$, we also have

$$f(y_n) = f'(x^*)\left(e_{n,y} + c_2 e_{n,y}^2 + c_3 e_{n,y}^3 + \ldots + c_8 e_{n,y}^8\right) + O(e_{n,y}^9). \tag{2.13}$$

According to the proof of Theorem 1, we have

$$\begin{aligned} e_{n,z} = z_n - x^* = {} & (-c_2c_3)e_n^4 + (20c_2^4 + 2c_2^2c_3 - 2c_3^2 - 2c_2c_4)e_n^5 + (-218c_2^5 + 156c_2^3c_3 \\ & + 3c_2^2c_4 - 7c_3c_4 + c_2(6c_3^2 - 3c_5))e_n^6 + 2(730c_2^6 - 1006c_2^4c_3 + 2c_3^3 \\ & + 118c_2^3c_4 + 8c_2c_3c_4 - 3c_4^2 - 5c_3c_5 + 2c_2^2(115c_3^2 - c_5))e_n^7 \\ & + (-7705c_2^7 + 15424c_2^5c_3 - 2946c_2^4c_4 + 1393c_2^2c_3c_4 + c_4(14c_3^2 - 17c_5) \\ & + 35c_2^3(-211c_3^2 + 9c_5) + 5c_2(121c_3^3 + 2c_4^2 + 4c_3c_5))e_n^8 + O(e_n^9). \end{aligned}$$

Also for $f(z_n)$, we get

$$f(z_n) = f^{'}(x^*)\left(e_{n,z} + c_2e_{n,z}^2 + c_3e_{n,z}^3 + \ldots + c_8e_{n,z}^8\right) + O(e_{n,z}^9). \tag{2.14}$$

By (2.11) and (2.13), we obtain

$$\begin{aligned} t_n = \frac{f(y_n)}{f(x_n)} &= c_2e_n + (-3c_2^2 + 2c_3)e_n^2 + (8c_2^3 - 10c_2c_3 + 3c_4)e_n^3 + (-20c_2^4 + 37c_2^2c_3 \\ &- 8c_3^2 - 14c_2c_4 + 4c_5)e_n^4 + (48c_2^5 - 118c_2^3c_3 + 51c_2^2c_4 - 22c_3c_4 + c_2(55c_3^2 \\ &- 18c_5))e_n^5 + (-112c_2^6 + 344c_2^4c_3 + 26c_3^3 - 163c_2^3c_4 + 150c_2c_3c_4 - 15c_4^2 \\ &- 28c_3c_5 + c_2^2(-252c_3^2 - 65c_5))e_n^6 + O(e_n^7). \end{aligned} \tag{2.15}$$

By (2.13) and (2.14), we obtain

$$\begin{aligned} s_n = \frac{f(z_n)}{f(y_n)} &= -c_3e_n^2 + (20c_2^3 - 2c_4)e_n^3 + (-178c_2^4 + 121c_2^2c_3 - c_3^2 - c_2c_4)e_n^4 + (1004c_2^5 \\ &- 1286c_2^3c_3 + 184c_2^2c_4 - 6c_3c_4 + c_2(240c_3^2 - 2c_5))e_n^5 + O(e_n^6). \end{aligned} \tag{2.16}$$

By (2.11) and (2.14), we get

$$\begin{aligned} u_n = \frac{f(z_n)}{f(x_n)} &= -c_2c_3e_n^3 + (20c_2^4 + 3c_2^2c_3 - 2c_3^2 - 2c_2c_4)e_n^4 + (-238c_2^5 + 153c_2^3c_3 + 5c_2^2c_4 \\ &- 7c_3c_4 + c_2(9c_3^2 - 3c_5))e_n^5 + O(e_n^6). \end{aligned} \tag{2.17}$$

Expanding H at $(0,0,0)$ yields

$$\begin{aligned} H(t_n, s_n, u_n) &= H_{0,0,0} + t_nH_{1,0,0} + s_nH_{0,1,0} + u_nH_{0,0,1} + t_ns_nH_{1,1,0} + t_nu_nH_{1,0,1} \\ &+ s_nu_nH_{0,1,1} + t_ns_nu_nH_{1,1,1} + \frac{t_n^2}{2}H_{2,0,0} + O(t_n^3, s_n^2, u_n^2). \end{aligned} \tag{2.18}$$

By substituting (2.11)-(2.18) into (2.10), we obtain

$$e_{n+1} = x_{n+1} - x^* = R_4e_n^4 + R_5e_n^5 + R_6e_n^6 + R_7e_n^7 + R_8e_n^8 + O(e_n^9),$$

where

$$\begin{aligned} R_4 &= \frac{1}{2}c_2\left(2c_3 + c_2^2(-10 + G_2)\right)(-1 + H_{0,0,0}), \\ R_5 &= c_2^2c_3\left(-2 + H_{1,0,0}\right), \\ R_6 &= \frac{1}{2}c_2c_3\left(-2c_3\left(-1 + H_{0,1,0}\right) + c_2^2\left(-12 + H_{2,0,0}\right)\right), \\ R_7 &= \frac{-1}{6}c_2^2c_3\left(c_2^2\left(36 + G_3\right) + 6c_3\left(-4 + H_{0,0,1} + H_{1,1,0}\right)\right), \\ R_8 &= c_2c_3\left(12c_2^4 + 14c_2^2c_3 - c_3^2 - c_2c_4\right). \end{aligned}$$

Clearly $R_8 \neq 0$, by setting $R_4 = R_5 = R_6 = R_7 = 0$, the convergence order becomes eight. Sufficient conditions are given by the following set of equations

$$\begin{aligned}
&H_{0,0,0} = 1, \quad G_2 = 10, && \Rightarrow \quad R_4 = 0,\\
&H_{1,0,0} = 2, && \Rightarrow \quad R_5 = 0,\\
&H_{0,1,0} = 1, \quad H_{2,0,0} = 12, && \Rightarrow \quad R_6 = 0,\\
&G_3 = -36, \quad H_{0,0,1} = 4, \quad H_{1,1,0} = 0, && \Rightarrow \quad R_7 = 0,
\end{aligned}$$

and the error equation becomes

$$e_{n+1} = R_8 e_n^8 + O(e_n^9),$$

which finishes the proof. □

2.3 Main contribution: Construction of an optimal four-point class

This section contains the main contribution. To increase the order of convergence, we add one Newton's step to the method (2.10) to get

$$\begin{cases}
y_n = x_n - \dfrac{f(x_n)}{f'(x_n)},\\
z_n = y_n - G(t_n) \cdot \dfrac{f(y_n)}{f'(x_n)},\\
w_n = z_n - H(t_n, s_n, u_n) \cdot \dfrac{f(z_n)}{f'(x_n)},\\
x_{n+1} = w_n - \dfrac{f(w_n)}{f'(w_n)}.
\end{cases} \tag{2.19}$$

The method (2.19) uses six function evaluations therefore the method is not an optimal four-point method. We modify (2.19) by approximating $f'(w_n)$ by

$$f'(w_n) \approx \frac{f'(x_n)}{I(t_n) + J(s_n) + K(u_n) + L(t_n, u_n) + M(p_n, q_n, r_n) + N(t_n, s_n, u_n, r_n)},$$

using only the values $f(x_n)$, $f(y_n)$, $f(z_n)$, $f(w_n)$ and $f'(x_n)$. More precisely, we use the abbreviations $t_n = \frac{f(y_n)}{f(x_n)}$, $s_n = \frac{f(z_n)}{f(y_n)}$, $u_n = \frac{f(z_n)}{f(x_n)}$, $p_n = \frac{f(w_n)}{f(x_n)}$, $q_n = \frac{f(w_n)}{f(y_n)}$, $r_n = \frac{f(w_n)}{f(z_n)}$ and utilize Mathematica [14] to carefully choose the weight functions $I, J, K : \mathbb{R} \to \mathbb{R}$, $L : \mathbb{R}^2 \to \mathbb{R}$, $M : \mathbb{R}^3 \to \mathbb{R}$ and $N : \mathbb{R}^4 \to \mathbb{R}$ from a class of admissible functions such that the scheme

$$\begin{cases}
y_n = x_n - \dfrac{f(x_n)}{f'(x_n)},\\
z_n = y_n - G(t_n) \cdot \dfrac{f(y_n)}{f'(x_n)},\\
w_n = z_n - H(t_n, s_n, u_n) \cdot \dfrac{f(z_n)}{f'(x_n)},\\
x_{n+1} = w_n - (I(t_n) + J(s_n) + K(u_n) + L(t_n, u_n) + M(p_n, q_n, r_n) + N(t_n, s_n, u_n, r_n)) \cdot \dfrac{f(w_n)}{f'(x_n)},
\end{cases} \tag{2.20}$$

is of order 16.

Theorem 3. *Assume that $f : D \subset \mathbb{R} \to \mathbb{R}$ is 16 times continuously differentiable and has a simple zero $x^* \in D$ and let $I, J, K : \mathbb{R} \to \mathbb{R}$, $L : \mathbb{R}^2 \to \mathbb{R}$, $M : \mathbb{R}^3 \to \mathbb{R}$ and $N : \mathbb{R}^4 \to \mathbb{R}$ be sufficiently differentiable functions. If*

$$
\begin{aligned}
&I_0 = 0, \quad I_1 = 2, \quad I_2 = 12,\\
&J_0 = 0, \quad J_1 = 1, \quad J_2 = 0, \quad J_3 = -6\\
&K_0 = 1, \quad K_1 = 4, \quad K_2 = -8,\\
&L_{0,0} = 0, \quad L_{1,0} = 0, \quad L_{1,1} = 1, \quad L_{2,0} = 0, \quad L_{0,1} = 0,\\
&L_{3,0} = 0, \quad L_{2,1} = 12, \quad L_{3,1} = 12, \quad L_{0,2} = 0, \quad L_{1,2} = -20,\\
&H_{0,1,1} = 0, \quad H_{1,1,1} = 0,\\
&M_{0,0,0} = 0, \quad M_{1,0,0} = 8, \quad M_{0,1,0} = 2, \quad M_{0,0,1} = 1,\\
&N_{0,0,0,0} = N_{1,0,0,0} = N_{0,1,0,0} = N_{0,0,1,0} = N_{0,0,0,1} = N_{2,0,0,0} = N_{1,1,0,0}\\
&\qquad = N_{0,2,0,0} = N_{1,0,1,0} = N_{2,1,0,0} = N_{3,1,0,0} = N_{2,2,0,0} = N_{0,1,0,1}\\
&\qquad = N_{3,0,0,0} = N_{4,0,0,0} = N_{3,0,1,0} = N_{1,1,0,1} = N_{2,1,1,0} = N_{3,0,0,1}\\
&\qquad = N_{0,0,1,1} = N_{3,2,0,0} = N_{4,1,0,0} = N_{1,2,0,0} = N_{2,0,1,0} = N_{1,1,1,0} = 0,\\
&N_{1,0,0,1} = 2, \quad N_{2,0,0,1} = 12, \quad N_{0,2,1,0} = -8, \quad N_{4,0,1,0} = 576, \quad N_{0,1,1,0} = 2.
\end{aligned}
$$

where $I_i = \frac{d^i I(t_n)}{dt_n^i}|_0$, $J_i = \frac{d^i J(s_n)}{ds_n^i}|_0$ and $K_i = \frac{d^i K(u_n)}{du_n^i}|_0$, $L_{i,j} = \frac{\partial^{i+j} L(t_n,u_n)}{\partial t_n^i \partial u_n^j}|_{(0,0)}$, $M_{i,j,k} = \frac{\partial^{i+j+k} M(p_n,q_n,r_n)}{\partial p_n^i \partial q_n^j \partial r_n^k}|_{(0,0,0)}$ and $N_{i,j,k,l} = \frac{\partial^{i+j+k+l} N(t_n,s_n,u_n,r_n)}{\partial t_n^i \partial s_n^j \partial u_n^k \partial r_n^l}|_{(0,0,0,0)}$ for $i, j, k, l = 0, 1, 2, 3, \ldots$, and the initial point x_0 is sufficiently close to x^*, then the sequence $\{x_n\}$ defined by (2.20) converges to x^* and the order of convergence is 16.

Proof. Let $e_n := x_n - x^*$, $e_{n,y} := y_n - x^*$, $e_{n,z} := z_n - x^*$, $e_{n,w} := w_n - x^*$ and $c_n := \frac{f^{(n)}(x^*)}{n! f'(x^*)}$ for $n \in \mathbb{N}$. Using the fact that $f(x^*) = 0$, Taylor's expansion of f at x^* yields

$$f(x_n) = f'(x^*)(e_n + c_2 e_n^2 + c_3 e_n^3 + \ldots + c_{16} e_n^{16}) + O(e_n^{17}), \tag{2.21}$$

and expanding f' at x^* we get

$$f'(x_n) = f'(x^*)(1 + 2c_2 e_n + 3c_3 e_n^2 + \ldots + 17c_{17} e_n^{16}) + O(e_n^{17}). \tag{2.22}$$

Therefore

$$
\begin{aligned}
\frac{f(x_n)}{f'(x_n)} &= e_n - c_2 e_n^2 + 2(c_2^2 - c_3)e_n^3 + (-4c_2^3 + 7c_2c_3 - 3c_4)e_n^4 + (8c_2^4 - 20c_2^2c_3 + 6c_3^2\\
&\quad + 10c_2c_4 - 4c_5)e_n^5 + 2(-16c_2^5 + 52c_2^3c_3 - 28c_2^2c_4 + 17c_3c_4 + c_2(-33c_3^2 + 13c_5))e_n^6\\
&\quad + \ldots + (-64c_2^7 + 304c_2^5c_3 - 176c_2^4c_4 + 348c_2^2c_3c_4 + c_2^3(-408c_3^2 + 92c_5)\\
&\quad + c_4(-75c_3^2 + 31c_5) + c_2(135c_3^3 - 64c_4^2 - 118c_3c_5))e_n^8 + O(e_n^9),
\end{aligned}
$$

so,

$$
\begin{aligned}
e_{n,y} = y_n - x^* &= c_2 e_n^2 + (-2c_2^2 + 2c_3)e_n^3 + (4c_2^3 - 7c_2c_3 + 3c_4)e_n^4 + (-8c_2^4 + 20c_2^2c_3 - 6c_3^2\\
&\quad - 10c_2c_4 + 4c_5)e_n^5 + 2(16c_2^5 - 52c_2^3c_3 + 28c_2^2c_4 - 17c_3c_4 + c_2(33c_3^2 - 13c_5))e_n^6\\
&\quad + \ldots + (64c_2^7 - 304c_2^5c_3 + 176c_2^4c_4 - 348c_2^2c_3c_4 + c_2^3(408c_3^2 - 92c_5)\\
&\quad + c_4(75c_3^2 - 31c_5) + c_2(-135c_3^3 + 64c_4^2 + 118c_3c_5))e_n^8 + O(e_n^9).
\end{aligned}
$$

For $f(y_n)$, we also have

$$f(y_n) = f^{'}(x^*)\left(e_{n,y} + c_2e_{n,y}^2 + c_3e_{n,y}^3 + \ldots + c_{16}e_{n,y}^{16}\right) + O(e_{n,y}^{17}). \tag{2.23}$$

According to the proof of Theorem 2, we have

$$\begin{aligned} e_{n,z} = z_n - x^* = {} & (-c_2c_3)e_n^4 + (20c_2^4 + 2c_2^2c_3 - 2c_3^2 - 2c_2c_4)e_n^5 + (-218c_2^5 + 156c_2^3c_3 \\ & + 3c_2^2c_4 - 7c_3c_4 + c_2(6c_3^2 - 3c_5))e_n^6 + 2(730c_2^6 - 1006c_2^4c_3 + 2c_3^3 \\ & + 118c_2^3c_4 + 8c_2c_3c_4 - 3c_4^2 - 5c_3c_5 + 2c_2^2(115c_3^2 - c_5))e_n^7 \\ & + (-7705c_2^7 + 15424c_2^5c_3 - 2946c_2^4c_4 + 1393c_2^2c_3c_4 + c_4(14c_3^2 - 17c_5) \\ & + 35c_2^3(-211c_3^2 + 9c_5) + 5c_2(121c_3^3 + 2c_4^2 + 4c_3c_5))e_n^8 + O(e_n^9). \end{aligned}$$

For $f(z_n)$, we also get

$$f(z_n) = f^{'}(x^*)\left(e_{n,z} + c_2e_{n,z}^2 + c_3e_{n,z}^3 + \ldots + c_{16}e_{n,z}^{16}\right) + O(e_{n,z}^{17}). \tag{2.24}$$

By (2.21) and (2.23), we obtain

$$\begin{aligned} t_n = \frac{f(y_n)}{f(x_n)} = {} & c_2e_n + (-3c_2^2 + 2c_3)e_n^2 + (8c_2^3 - 10c_2c_3 + 3c_4)e_n^3 + (-20c_2^4 + 37c_2^2c_3 \\ & - 8c_3^2 - 14c_2c_4 + 4c_5)e_n^4 + (48c_2^5 - 118c_2^3c_3 + 51c_2^2c_4 - 22c_3c_4 + c_2(55c_3^2 \\ & - 18c_5))e_n^5 + (-112c_2^6 + 344c_2^4c_3 + 26c_3^3 - 163c_2^3c_4 + 150c_2c_3c_4 - 15c_4^2 \\ & - 28c_3c_5 + c_2^2(-252c_3^2 + 65c_5))e_n^6 + (256c_2^7 - 944c_2^5c_3 + 480c_2^4c_4 - 693c_2^2c_3c_4 \\ & + c_2^3(952c_3^2 - 207c_5) + c_4(105c_3^2 - 38c_5) + 2c_2(-114c_3^3 + 51c_4^2 + 95c_3c_5))e_n^7 \\ & + (-576c_2^8 + 2480c_2^6c_3 - 1336c_2^5c_4 + 2660c_2^3c_3c_4 + 6c_2c_4(-156c_3^2 + 43c_5) \\ & + c_2^4(-3200c_3^2 + 607c_5) + 3c_2^2(418c_3^3 - 159c_4^2 - 292c_3c_5) + 3(-24c_3^4 + 47c_3c_4^2 \\ & + 44c_3^2c_5 - 8c_5))e_n^8 + O(e_n^9). \end{aligned} \tag{2.25}$$

By (2.23) and (2.24), we obtain

$$\begin{aligned} s_n = \frac{f(z_n)}{f(y_n)} = {} & -c_3e_n^2 + (20c_2^3 - 2c_4)e_n^3 + (-178c_2^4 + 121c_2^2c_3 - c_3^2 - c_2c_4 - 3c_5)e_n^4 + (1004c_2^5 \\ & - 1286c_2^3c_3 + 184c_2^2c_4 - 6c_3c_4 + c_2(240c_3^2 - 2c_5))e_n^5 + (-4567c_2^6 + 8541c_2^4c_3 \\ & + 155c_3^3 + 1863c_2^3c_4 + 725c_2c_3c_4 - 7c_4^2 - 10c_3c_5 + \frac{1}{2}c_2^2(-6866c_3^2 + 492c_5))e_n^6 \\ & + \frac{1}{6}(109500c_2^7 - 269472c_2^5c_3 + 72756c_2^4c_4 - 59376c_2^2c_3c_4 + c_2^3(172248c_3^2 - 14616c_5) \\ & + 12c_4(348c_3^2 - 11c_5) + c_2(-24048c_3^3 + 3276c_4^2 + 5796c_3c_5))e_n^7 + (-66359c_2^8 \\ & + 204034c_2^6c_3 - 1725c_3^4 - 63159c_2^5c_4 + 81203c_2^3c_3c_4 + 1038c_3c_4^2 + 923c_3^2c_5 - 17c_5^2 \\ & + c_2c_4(-17234c_3^2 + 1453c_5) + c_2^4(-181979c_3^2 + 15700c_5) + 2c_2^2(23804c_3^3 - 3559c_4^2 \\ & - 6452c_3c_5))e_n^8 + O(e_n^9). \end{aligned} \tag{2.26}$$

By (2.21) and (2.24), we have

$$\begin{aligned} u_n = \frac{f(z_n)}{f(x_n)} &= -c_2c_3e_n^3 + (20c_2^4 + 3c_2^2c_3 - 2c_3^2 - 2c_2c_4)e_n^4 + (-238c_2^5 + 153c_2^3c_3 + 5c_2^2c_4 \\ &\quad - 7c_3c_4 + c_2(9c_3^2 - 3c_5))e_n^5 + (1698c_2^6 - 2185c_2^4c_3 + 6c_3^3 + 231c_2^3c_4 + 26c_2c_3c_4 \\ &\quad - 6c_4^2 - 10c_3c_5 + 7c_2^2(64c_3^2 + c_5))e_n^6 + (-9403c_2^7 + 17847c_2^5c_3 - 3197c_2^4c_4 \\ &\quad + 1359c_2^2c_3c_4 + c_4(23c_3^2 - 17c_5) + c_2^3(-7985c_3^2 + 308c_5) + 2c_2(295c_3^3 + 9c_4^2 \\ &\quad + 17c_3c_5))e_n^7 + (44503c_2^8 - 111251c_2^6c_3 + 292c_3^4 + 25635c_2^5c_4 - 23315c_2^3c_3c_4 \\ &\quad + 27c_3c_4^2 + 382c_2^4(203c_3^2 - 11c_5) + 28c_3^2c_5 - 12c_5^2 + 2c_2c_4(1344c_3^2 + 23c_5) \\ &\quad + c_2^2(-14492c_3^3 + 1031c_4^2 + 1816c_3c_5))e_n^8 + O(e_n^9), \end{aligned} \tag{2.27}$$

According to the proof of Theorem 2, we have

$$\begin{aligned} e_{n,w} = w_n - x^* &= c_2c_3(12c_2^4 + 14c_2^2c_3 - c_3^2 - c_2c_4)e_n^8 + 2(120c_2^8 + 344c_2^6c_3 - 37c_2^4c_3^2 + c_3^4 \\ &\quad - 22c_2^5c_4 - 30c_2^3c_3c_4 + 5c_2c_3^2c_4 + c_2^2(-40c_3^3 + c_4^2 + c_3c_5))e_n^9 + (10296c_2^9 + 5059c_2^7c_3 \\ &\quad - 1622c_2^6c_4 + 452c_2^4c_3c_4 - 19c_3^3c_4 + c_2^2c_4(456c_3^2 - 7c_5) + c_2^3(236c_3^3 + 63c_4^2 \\ &\quad + 91c_3c_5))e_n^{10} + O(e_n^{11}). \end{aligned} \tag{2.28}$$

For $f(w_n)$, we also obtain

$$f(w_n) = f^{'}(x^*)\left(e_{n,w} + c_2e_{n,w}^2 + c_3e_{n,w}^3 + \ldots + c_{16}e_{n,w}^{16}\right) + O(e_{n,w}^{17}). \tag{2.29}$$

By (2.21) and (2.29), we have

$$\begin{aligned} p_n = \frac{f(w_n)}{f(x_n)} &= c_2c_3(12c_2^4 + 14c_2^2c_3 - c_3^2 - c_2c_4)e_n^7 + (-240c_2^8 - 700c_2^6c_3 + 60c_2^4c_3^2 - 2c_3^4 \\ &\quad + 44c_2^5c_4 + 61c_2^3c_3c_4 - 10c_2c_3^2c_4 + c_2^2(81c_3^3 - 2c_4^2 - 2c_3c_5))e_n^8 + (10536c_2^9 \\ &\quad + 5759c_2^7c_3 - 1666c_2^6c_4 + 391c_2^4c_3c_4 - 19c_3^3c_4 + c_2^2c_4(467c_3^2 - 7c_5) + c_2^5(-8121c_3^2 \\ &\quad + 76c_5) + c_2c_3(151c_3^3 - 26c_4^2 - 17c_3c_5) + c_2^3(141c_3^3 + 65c_4^2 + 93c_3c_5))e_n^9 + O(e_n^{10}). \end{aligned} \tag{2.30}$$

By (2.23) and (2.29), we get

$$\begin{aligned} q_n = \frac{f(w_n)}{f(y_n)} &= -c_3(-12c_2^4 - 14c_2^2c_3 + c_3^2 + c_2c_4)e_n^6 - 2(120c_2^7 + 332c_2^5c_3 - 39c_2^3c_3^2 - 22c_2^4c_4 \\ &\quad - 29c_2^2c_3c_4 + 4c_3^2c_4 + c_2(-25c_3^3 + c_4^2 + c_3c_5))e_n^7 + (9816c_2^8 + 4151c_2^6c_3 - 1534c_2^5c_4 \\ &\quad + 449c_2^3c_3c_4 + c_2c_4(275c_3^2 - 7c_5) + c_2^4(-6551c_3^2 + 76c_5) + c_3(41c_3^3 - 19c_4^2 - 13c_3c_5) \\ &\quad + c_2^2(288c_3^3 + 59c_4^2 + 87c_3c_5))e_n^8 + O(e_n^9). \end{aligned} \tag{2.31}$$

By (2.24) and (2.29), we obtain

$$\begin{aligned} r_n = \frac{f(w_n)}{f(z_n)} &= (-12c_2^4 - 14c_2^2c_3 + c_3^2 + c_2c_4)e_n^4 + 2(384c_2^5 - 58c_2^3c_3c_3 - 30c_2^2c_4 + 6c_3c_4 \\ &+ 2c_2(-25c_3^2 + c_5))e_n^5 + (-4271c_2^6 + 3691c_2^4c_3 - 42c_3^3 - 78c_2^3c_4 - 177c_2c_3c_4 + 7c_4^2 \\ &+ 10c_3c_5 - c_2^2(148c_3^2 + 45c_5))e_n^6 + 2(15680c_2^7 - 23481c_2^5c_3 + 2760c_2^4c_4 - 237c_2^2c_3c_4 \\ &+ c_2^3(7182c_3^2 - 51c_5) + c_4(-102c_3^2 + 11c_5) - c_2(126c_3^3 + 75c_4^2 + 125c_3c_5))e_n^7 \\ &+ (-182548c_2^8 + 384369c_2^6c_3 - 191c_3^4 - 68503c_2^5c_4 + 43033c_2^3c_3c_4 - 321c_3c_4^2 \\ &- 278c_3^2c_5 + 17c_5^2 - c_2c_4(1316c_3^2 + 417c_5) + c_2^4(-213934c_3^2 + 7347c_5) \\ &+ c_2^2(28040c_3^3 - 398c_4^2 - 673c_3c_5)e_n^8 + O(e_n^9). \end{aligned} \tag{2.32}$$

Expanding I, J, K, L, M and N at 0 in $\mathbb{R}, \mathbb{R}^2, \mathbb{R}^3$ and $\mathbb{R}^4$, respectively, yield

$$I(t_n) = I_0 + t_nI_1 + \frac{t_n^2}{2}I_2 + O(t_n^3), \tag{2.33}$$

$$J(s_n) = J_0 + s_nJ_1 + \frac{s_n^2}{2}J_2 + \frac{s_n^3}{6}J_3 + O(s_n^4), \tag{2.34}$$

$$K(u_n) = K_0 + u_nK_1 + \frac{u_n^2}{2}K_2 + O(u_n^3), \tag{2.35}$$

$$\begin{aligned} L(t_n, u_n) &= L_{0,0} + t_nL_{1,0} + u_nL_{0,1} + t_nu_nL_{1,1} + \frac{t_n^2}{2}L_{2,0} + \frac{u_n^2}{2}L_{0,2} + \frac{t_nu_n^2}{2}L_{1,2} \\ &+ \frac{t_n^2u_n}{2}L_{2,1} + \frac{t_n^2u_n^2}{4}L_{2,2} + \frac{t_n^3}{6}L_{3,0} + \frac{t_n^3u_n}{6}L_{3,1} + \frac{t_n^3u_n^2}{12}L_{3,2} + O(t_n^4, u_n^3), \end{aligned} \tag{2.36}$$

$$\begin{aligned} M(p_n, q_n, r_n) &= M_{0,0,0} + p_nM_{1,0,0} + q_nM_{0,1,0} + r_nM_{0,0,1} + p_nq_nM_{1,1,0} + p_nr_nM_{1,0,1} + p_nq_nr_nM_{1,1,1} \\ &+ q_nr_nM_{0,1,1} + \frac{p_n^2}{2}M_{2,0,0} + \frac{q_n^2}{2}M_{0,2,0} + \frac{r_n^2}{2}M_{0,0,2} + \frac{p_nr_n^2}{2}M_{1,0,2} + \frac{p_n^2r_n}{2}M_{2,0,1} \\ &+ \frac{p_n^2q_n}{2}M_{2,1,0} + \frac{p_nq_n^2}{2}M_{1,2,0} + \frac{p_nr_n^2}{2}M_{1,0,2} + \frac{q_nr_n^2}{2}M_{0,1,2} + \frac{q_n^2r_n}{2}M_{0,2,1} \\ &+ \frac{p_nq_nr_n^2}{2}M_{1,1,2} + \frac{p_nq_n^2r_n}{2}M_{1,2,1} + \frac{p_n^2q_nr_n}{2}M_{2,1,1} + \frac{q_n^2r_n^2}{4}M_{0,2,2} + \frac{p_nq_n^2r_n^2}{4}M_{1,2,2} \\ &+ \frac{p_n^2r_n^2}{4}M_{2,0,2} + \frac{p_n^2q_nr_n^2}{4}M_{2,1,2} + \frac{p_n^2q_n^2}{4}M_{2,2,0} + \frac{p_n^2q_n^2r_n}{4}M_{2,2,1} + \frac{p_n^2q_n^2r_n^2}{8}M_{2,2,2} \\ &+ O(p_n^3, q_n^3, r_n^3), \end{aligned} \tag{2.37}$$

and,

$$
\begin{aligned}
N(t_n, s_n, u_n, r_n) &= N_{0,0,0,0} + t_n N_{1,0,0,0} + s_n N_{0,1,0,0} + u_n N_{0,0,1,0} + r_n N_{0,0,0,1} + t_n s_n N_{1,1,0,0} \\
&+ t_n u_n N_{1,0,1,0} + t_n r_n N_{1,0,0,1} + u_n r_n N_{0,0,1,1} + s_n u_n N_{0,1,1,0} + s_n r_n N_{0,1,0,1} \\
&+ t_n s_n u_n N_{1,1,1,0} + t_n s_n r_n N_{1,1,0,1} + s_n u_n r_n N_{0,1,1,1} + t_n u_n r_n N_{1,0,1,1} \\
&+ t_n s_n u_n r_n N_{1,1,1,1} + \frac{t_n^2}{2} N_{2,0,0,0} + \frac{s_n^2}{2} N_{0,2,0,0} + \frac{s_n^2 r_n}{2} N_{0,2,0,1} + \frac{s_n^2 u_n}{2} N_{0,2,1,0} \\
&+ \frac{s_n^2 u_n r_n}{2} N_{0,2,1,1} + \frac{t_n s_n^2}{2} N_{1,2,0,0} + \frac{t_n s_n^2 r_n}{2} N_{1,2,0,1} + \frac{t_n s_n^2 u_n}{2} N_{1,2,1,0} \\
&+ \frac{t_n^2 r_n}{2} N_{2,0,0,1} + \frac{t_n^2 u_n}{2} N_{2,0,1,0} + \frac{t_n^2 s_n}{2} N_{2,1,0,0} + \frac{t_n^2 u_n r_n}{2} N_{2,0,1,1} \\
&+ \frac{t_n^2 s_n}{2} N_{2,1,0,0} + \frac{t_n^2 s_n r_n}{2} N_{2,1,0,1} + \frac{t_n^2 s_n u_n}{2} N_{2,1,1,0} + \frac{t_n^2 s_n u_n r_n}{2} N_{2,1,1,1} \\
&+ \frac{t_n^2 s_n^2}{4} N_{2,2,0,0} + \frac{t_n^2 s_n^2 r_n}{4} N_{2,2,0,1} + \frac{t_n^2 s_n^2 u_n}{4} N_{2,2,1,0} + \frac{t_n^2 s_n^2 u_n r_n}{4} N_{2,2,1,1} \\
&+ \frac{t_n^3}{6} N_{3,0,0,0} + \frac{t_n^3 r_n}{6} N_{3,0,0,1} + \frac{t_n^3 u_n}{6} N_{3,0,1,0} + \frac{t_n^3 u_n r_n}{6} N_{3,0,1,1} \\
&+ \frac{t_n^3 s_n}{6} N_{3,1,0,0} + \frac{t_n^3 s_n r_n}{6} N_{3,1,0,1} + \frac{t_n^3 s_n u_n}{6} N_{3,1,1,0} + \frac{t_n^3 s_n u_n r_n}{6} N_{3,1,1,1} \\
&+ \frac{t_n^3 s_n^2}{12} N_{3,2,0,0} + \frac{t_n^3 s_n^2 r_n}{12} N_{3,2,0,1} + \frac{t_n^3 s_n^2 u_n}{12} N_{3,2,1,0} + \frac{t_n^3 s_n^2 u_n r_n}{12} N_{3,2,1,1} \\
&+ \frac{t_n^4}{24} N_{4,0,0,0} + \frac{t_n^4 r_n}{24} N_{4,0,0,1} + \frac{t_n^4 u_n}{24} N_{4,0,1,0} + \frac{t_n^4 s_n r_n}{24} N_{4,1,0,1} + \frac{t_n^4 s_n}{24} N_{4,1,0,0} \\
&+ \frac{t_n^4 s_n r_n}{24} N_{4,1,0,1} + \frac{t_n^4 s_n u_n}{24} N_{4,1,1,0} + \frac{t_n^4 s_n u_n r_n}{24} N_{4,1,1,1} + \frac{t_n^4 s_n^2}{48} N_{4,2,0,0} \\
&+ \frac{t_n^4 s_n^2 r_n}{48} N_{4,2,0,1} + \frac{t_n^4 s_n^2 u_n}{48} N_{4,2,1,0} + \frac{t_n^4 s_n^2 u_n r_n}{48} N_{4,2,1,1} + O(t_n^5, s_n^3, u_n^2, r_n^2).
\end{aligned} \tag{2.38}
$$

By substituting (2.21)-(2.38) into (2.20), we obtain

$$
\begin{aligned}
e_{n+1} = x_{n+1} - x^* &= R_8 e_n^8 + R_9 e_n^9 + R_{10} e_n^{10} + R_{11} e_n^{11} + R_{12} e_n^{12} \\
&+ R_{13} e_n^{13} + R_{14} e_n^{14} + R_{15} e_n^{15} + R_{16} e_n^{16} + O(e_n^{17}),
\end{aligned}
$$

where

$$
R_8 = -c_2 c_3 \left(12c_2^4 + 14c_2^2 c_3 - c_3^2 - c_2 c_4\right) \left(-1 + I_0 + J_0 + K_0 + L_{0,0} + M_{0,0,0} + N_{0,0,0,0}\right)
$$

$$
R_9 = c_2^2 c_3 \left(-12c_2^4 - 14c_2^2 c_3 + c_3^2 + c_2 c_4\right) \left(-2 + I_1 + L_{1,0} + N_{1,0,0,0}\right),
$$

$$
\begin{aligned}
R_{10} = &\frac{-1}{2} c_2 c_3 \left(12c_2^4 + 14c_2^2 c_3 - c_3^2 - c_2 c_4\right) \\
&\left(-2c_3 \left(-1 + J_1 + N_{0,1,0,0}\right) + c_2^2(-12 + I_2 + L_{2,0} + N_{2,0,0,0})\right),
\end{aligned}
$$

$$R_{11} = \frac{-1}{6} c_2^2 c_3 \left(12c_2^4 + 14c_2^2 c_3 - c_3^2 - c_2 c_4\right) \big(-6c_3 \left(-4 + K_1 + L_{0,1} + N_{0,0,1,0} + N_{1,1,0,0}\right) + c_2^2 \left(L_{3,0} + N_{3,0,0,0}\right) \big),$$

$$R_{12} = \frac{1}{24} c_2 c_3 \left(12c_2^4 + 14c_2^2 c_3 - c_3^2 - c_2 c_4\right) (-24c_2 c_4(-1 + M_{0,0,1} + N_{0,0,0,1}) - 12c_3^2(J_2 + 2(-1 + M_{0,0,1} + N_{0,0,0,1}) + N_{0,2,0,0}) + 12c_2^2 c_3(2(-15 + L_{1,1} + 14M_{0,0,1} + 14N_{0,0,0,1} + N_{1,0,1,0}) + N_{2,1,0,0}) + c_2^4(288(-1 + M_{0,0,1} + N_{0,0,0,1}) - N_{4,0,0,0})),$$

$$R_{13} = \frac{1}{6} c_2^2 c_3 (12c_2^4 + 14c_2^2 c_3 - c_3^2 - c_2 c_4) (72c_2^4(-2 + N_{1,0,0,1}) - 6c_2 c_4(-2 + N_{1,0,0,1}) + 3c_3^2(2H_{0,1,1} - 2(-4 + N_{0,1,1,0} + N_{1,0,0,1}) - N_{1,2,0,0}) + c_2^2 c_3(3(-68 + L_{2,1} + 28N_{1,0,0,1} + N_{2,0,1,0}) + N_{3,1,0,0})),$$

$$R_{14} = \frac{1}{24} c_2 c_3 (12c_2^4 + 14c_2^2 c_3 - c_3^2 - c_2 c_4) (24c_2 c_3 c_4(-2 + M_{0,1,0} + N_{0,1,0,1}) + 4c_3^3(J_3 + 6(-1 + M_{0,1,0} + N_{0,1,0,1})) + 144c_2^6(-12 + N_{2,0,0,1}) - 12c_2^3 c_4(-12 + N_{2,0,0,1}) - 6c_2^2 c_3^2(2(-60 + K_2 + L_{0,2} - 2H_{1,1,1} + 28M_{0,1,0} + 28N_{0,1,0,1} + 2N_{1,1,1,0} + N_{2,0,0,1}) + N_{2,2,0,0}) + c_2^4 c_3(4(-372 + L_{3,1} - 72M_{0,1,0} - 72N_{0,1,0,1} + 42N_{2,0,0,1} + N_{3,0,1,0}) + N_{4,1,0,0})),$$

$$R_{15} = \frac{1}{24} c_2^2 c_3 (12c_2^4 + 14c_2^2 c_3 - c_3^2 - c_2 c_4) (24c_2 c_3 c_4(-8 + M_{1,0,0} + N_{0,0,1,1} + N_{1,1,0,1}) + 12c_3^3(-8 + 2M_{1,0,0} + 2N_{0,0,1,1} + N_{0,2,1,0} + 2N_{1,1,0,1}) + 48c_2^6 N_{3,0,0,1} - 4c_2^3 c_4 N_{3,0,0,1} - 2c_2^2 c_3^2(6(L_{1,2} + 4(-51 + 7M_{1,0,0} + 7N_{0,0,1,1} + 7N_{1,1,0,1}) + N_{2,1,1,0} + 2N_{3,0,0,1} + N_{3,2,0,0}) + c_2^4 c_3(-288(-6 + M_{1,0,0} + N_{0,0,1,1} + N_{1,1,0,1}) + 56N_{3,0,0,1} + N_{4,0,1,0})),$$

$$R_{16} = \frac{-1}{48} c_2 c_3 (12c_2^4 + 14c_2^2 c_3 - c_3^2 - c_2 c_4) (24c_3^4(-2 + M_{0,0,2} + N_{0,2,0,1}) + 24c_2 c_3^2 c_4(-6 + 2M_{0,0,2} + N_{0,2,0,1}) - 24c_2^3 c_3 c_4(-88 + 28M_{0,0,2} + 2N_{1,0,1,1} + N_{2,1,0,1}) - 24c_2^2(2c_3 c_5 - c_4^2(-2 + M_{0,0,2}) + c_3^3(-86 + 28M_{0,0,2} + 14N_{0,2,0,1} + 2N_{1,0,1,1} + N_{1,2,1,0} + 6N_{1,2,1,0} + N_{2,1,0,1})) + 4c_2^6 c_3(36(-145 + 56M_{0,0,2} + 4N_{1,0,1,1} + 2N_{2,1,0,1}) - 7N_{4,0,0,1}) + 24c_2^8(144(-2 + M_{0,0,2}) - N_{4,0,0,1}) + 2c_2^5 c_4(576 - 288M_{0,0,2} + N_{4,0,0,1}) + c_2^4 c_3^2 (2(6L_{2,2} + 4(6(86M_{0,0,2} - 6N_{0,2,0,1} + 7(-45 + 2N_{1,0,1,1} + N_{2,1,0,1})) + N_{3,1,1,0}) + N_{4,0,0,1}) + N_{4,2,0,0})).$$

In general $R_{16} \neq 0$, however, by setting $R_8 = R_9 = R_{10} = R_{11} = R_{12} = R_{13} = R_{15} = R_{15} = 0$, the

convergence order becomes 16. Sufficient conditions are given by the following set of equations

$$K_0 = 1, \quad I_0 = J_0 = L_{0,0} = M_{0,0,0} = N_{0,0,0,0} = 0, \qquad \Rightarrow \quad R_8 = 0,$$

$$I_1 = 2, \quad L_{1,0} = N_{1,0,0,0} = 0, \qquad \Rightarrow \quad R_9 = 0,$$

$$I_2 = 12, \quad J_1 = 1, \quad L_{2,0} = N_{0,1,0,0} = N_{2,0,0,0} = 0, \qquad \Rightarrow \quad R_{10} = 0,$$

$$K_1 = 4, \quad L_{0,1} = L_{3,0} = N_{0,0,1,0} = N_{1,1,0,0} = N_{3,0,0,0} = 0, \qquad \Rightarrow \quad R_{11} = 0,$$

$$L_{1,1} = 1, \quad M_{0,0,1} = 1, \quad J_2 = N_{0,0,0,1} = N_{0,2,0,0} = N_{4,0,0,0} = N_{1,0,1,0} = N_{2,1,0,0} = 0, \qquad \Rightarrow \quad R_{12} = 0,$$

$$L_{2,1} = 12, \quad N_{1,0,0,1} = N_{0,1,1,0} = 2, \quad H_{0,1,1} = N_{1,2,0,0} = N_{2,0,1,0} = N_{3,1,0,0} = 0, \qquad \Rightarrow \quad R_{13} = 0,$$

$$\begin{aligned} &J_3 = -6, \quad K_2 = -8, \quad L_{3,1} = 12, \quad M_{0,1,0} = 2, \quad N_{2,0,0,1} = 12, \\ &L_{0,2} = H_{1,1,1} = N_{0,1,0,1} = N_{1,1,1,0} = N_{2,2,0,0} = N_{3,0,1,0} = N_{4,1,0,0} = 0 \qquad \Rightarrow \quad R_{14} = 0, \end{aligned}$$

$$\begin{aligned} &L_{1,2} = -20, \quad M_{1,0,0} = 8, \quad N_{0,2,1,0} = -8 \quad N_{4,0,1,0} = 576, \\ &N_{0,0,1,1} = N_{1,1,0,1} = N_{3,0,0,1} = N_{2,1,1,0} = N_{3,2,0,0} = 0 \qquad \Rightarrow \quad R_{15} = 0, \end{aligned}$$

$$\begin{aligned} &L_{2,2} = 12, \quad M_{0,0,2} = 2, \quad N_{1,0,1,1} = 30 \quad N_{1,2,1,0} = -18, \\ &N_{0,2,0,1} = N_{2,1,0,1} = N_{4,0,0,1} = N_{3,1,1,0} = N_{4,2,0,0} = 0 \qquad \Rightarrow \quad R_{16} \neq 0, \end{aligned}$$

and the error equation becomes

$$e_{n+1} = \left(-c_2^2 c_3^2 (12c_2^4 - c_3^2 - c_2 c_4)(93c_2^5 - c_3 c_4 - c_2 c_5)\right) e_n^{16} + O(e_n^{17}),$$

which finishes the proof. □

3 Numerical results

3.1 Numerical implementation and comparison

In this section, three concrete methods of each of the families (2.10) and(2.20) are tested on a number of nonlinear equations. To obtain a high accuracy and avoid the loss of significant digits, we employed multi-precision arithmetic with 7000 significant decimal digits in the programming package of Mathematica 8.

Method 1. Weight functions G and H in (2.10) are given by

$$\begin{aligned} G(t_n) &= -6t_n^3 + 5t_n^2 + 2t_n + 1, \\ H(t_n, s_n, u_n) &= 1 + 2t_n + 4u_n + 6t_n^2 + s_n, \end{aligned} \tag{3.1}$$

where $t_n = \frac{f(y_n)}{f(x_n)}$, $s_n = \frac{f(z_n)}{f(y_n)}$ and $u_n = \frac{f(z_n)}{f(x_n)}$.

These functions satisfy the given conditions in Theorems 1 and 2, so

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - (-6t_n^3 + 5t_n^2 + 2t_n + 1) \cdot \dfrac{f(y_n)}{f'(x_n)}, \\ x_{n+1} = z_n - (1 + 2t_n + 4u_n + 6t_n^2 + s_n) \cdot \dfrac{f(z_n)}{f'(x_n)}. \end{cases} \tag{3.2}$$

Method 2. The method by H.T. Kung and J.F. Traub [18] is given by

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - G_f(x_n), \\ x_{n+1} = z_n - f^2(x_n) f(y_n) H_f(x_n, y_n, z_n), \end{cases} \tag{3.3}$$

where

$$\begin{aligned} G_f(x_n) &= \frac{f^2(x_n) f(y_n)}{f'(x_n)(f(x_n) - f(y_n))^2}, \\ H_f(x_n, y_n, z_n) &= G_f(x_n) \\ &\quad \Big(\frac{-1}{f^2(x_n)(f(x_n) - f(z_n))} + \frac{f(y_n) - f(x_n)}{f(x_n) f(y_n)(f(x_n) - f(z_n))^2} \\ &\quad + \frac{1}{(f(y_n) - f(z_n))(f(x_n) - f(z_n))^2}\Big). \end{aligned}$$

Method 3. The method by B. Neta [22] is given by

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \dfrac{f(x_n) + A f(y_n)}{f(x_n) + (A-2) f(y_n)} \dfrac{f(y_n)}{f'(x_n)}, \quad A \in \mathbb{R}, \\ x_{n+1} = y_n + \delta_1 f^2(x_n) + \delta_2 f^3(x_n), \end{cases} \tag{3.4}$$

where

$$\begin{gathered} F_y = f(y_n) - f(x_n), \quad F_z = f(z_n) - f(x_n), \\ \zeta_y = \frac{1}{F_y}\left(\frac{y_n - x_n}{F_y} - \frac{1}{f'(x_n)}\right), \zeta_z = \frac{1}{F_z}\left(\frac{z_n - x_n}{F_z} - \frac{1}{f'(x_n)}\right), \\ \delta_1 = \zeta_y + \delta_2 F_y, \qquad \delta_2 = -\frac{\zeta_y - \zeta_z}{F_y - F_z}, \end{gathered}$$

Method 4. The method by Khattri and Steihaug [16] is given by

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(y_n)}{\frac{x_n - y_n + \alpha f(x_n)}{(x_n - y_n)\alpha} - \frac{(x_n - y_n) f(x_n + \alpha f(x_n))}{(x_n - y_n + \alpha f(x_n))\alpha f(x_n)} - \frac{(2x_n - 2y_n + \alpha f(x_n)) f(y_n)}{(x_n - y_n)(x_n - y_n + \alpha f(x_n))}}, \quad \alpha \in \mathbb{R}, \\ x_{n+1} = z_n - \frac{f(z_n)}{H_1 f(x_n) + H_2 f(x_n + \alpha f(x_n)) + H_3 f(y_n) + H_4 f(z_n)}, \end{cases} \tag{3.5}$$

where

$$H_1 = \frac{(y_n - z_n)(z_n - x_n - \alpha f(x_n))}{\alpha f(x_n) + (y_n - x_n)(z_n - x_n)},$$
$$H_2 = \frac{-x_n y_n + x_n z_n + y_n z_n - z_n^2}{\alpha f(x_n)(-\alpha f(x_n) + y_n - x_n)(\alpha f(x_n) + x_n - z_n)},$$
$$H_3 = -\frac{(x_n + \alpha f(x_n))x_n - (x_n + \alpha f(x_n))z_n - x_n z_n + z_n^2}{(\alpha f(x_n) + x_n - y_n)(y_n - x_n)(y_n - z_n)},$$
$$H_4 = -\frac{(x_n + \alpha f(x_n))x_n + (x_n + \alpha f(x_n))y_n + x_n y_n - 2(x_n + \alpha f(x_n))z_n - 2x_n z_n - 2y_n z_n + 3z_n^2}{(\alpha f(x_n) + x_n - z_n)(z_n - x_n)(z_n - y_n)}.$$

Method 5. Weight functions G, H, I, J, K, L, M and N in (2.20) are given by

$$\begin{aligned}
&G(t_n) = -6t_n^3 + 5t_n^2 + 2t_n + 1,\\
&H(t_n, s_n, u_n) = 1 + 2t_n + 4u_n + 6t_n^2 + s_n,\\
&I(t_n) = 6t_n^2 + 2t_n, \quad J(s_n) = -s_n^3 + s_n + 1, \quad K(u_n) = 4u_n - 4u_n^2,\\
&L(t_n, u_n) = t_n u_n + 6t_n^2 u_n + 2t_n^3 u_n - 10t_n u_n^2, \quad M(p_n, q_n, r_n) = r_n + 2q_n + 8p_n,\\
&N(t_n, s_n, u_n, r_n) = 2t_n r_n + 2s_n u_n + 6t_n^2 r_n - 4s_n^2 u_n + 24t_n^4 u_n,
\end{aligned} \tag{3.6}$$

where $t_n = \frac{f(y_n)}{f(x_n)}$, $s_n = \frac{f(z_n)}{f(y_n)}$, $u_n = \frac{f(z_n)}{f(x_n)}$, $p_n = \frac{f(w_n)}{f(x_n)}$, $q_n = \frac{f(w_n)}{f(y_n)}$ and $r_n = \frac{f(w_n)}{f(z_n)}$.

These functions satisfy the given conditions in Theorems 1, 2, and 3, so

$$\begin{cases}
y_n = x_n - \dfrac{f(x_n)}{f'(x_n)},\\
z_n = y_n - (-6t_n^3 + 5t_n^2 + 2t_n + 1)\cdot\dfrac{f(y_n)}{f'(x_n)},\\
w_n = z_n - (1 + 2t_n + 4u_n + 6t_n^2 + s_n)\cdot\dfrac{f(z_n)}{f'(x_n)},\\
x_{n+1} = w_n - \big[1 + 6t_n^2 + 2t_n - s_n^3 + s_n + 4u_n - 4u_n^2\\
\qquad +t_n u_n + 6t_n^2 u_n + 2t_n^3 u_n - 10t_n u_n^2 + r_n + 2q_n + 8p_n\\
\qquad +2t_n r_n + 2s_n u_n + 6t_n^2 r_n - 4s_n^2 u_n + 24t_n^4 u_n\big]\cdot\dfrac{f(w_n)}{f'(x_n)}.
\end{cases} \tag{3.7}$$

Method 6. Weight functions G, H, I, J, K, L, M, and N in (2.20) are given by

$$\begin{aligned}
&G(t_n) = t_n^2(5 - 7t_n) + (2t_n + 1)(t_n^3 + 1) - 2t_n^4,\\
&H(t_n, s_n, u_n) = (1 + s_n) + (6 + u_n^2)(u_n + t_n^2) + 2(t_n - u_n),\\
&I(t_n) = (1 + t_n)(2t_n + t_n^2) + t_n^2(3 - t_n), \quad J(s_n) = \frac{s_n + s_n^2 - s_n^3}{1 + s_n}, \quad K(u_n) = \frac{1 + 5u_n}{1 + u_n},\\
&L(t_n, u_n) = t_n u_n + 6t_n^2 u_n + \frac{2t_n^3 u_n - 10t_n u_n^2}{1 + t_n u_n}, \quad M(p_n, q_n, r_n) = 2(p_n + q_n) + \frac{6p_n + r_n}{1 + p_n},\\
&N(t_n, s_n, u_n, r_n) = 8t_n^2 r_n - 4s_n^2 u_n - 2t_n^3 r_n + \frac{2s_n u_n + 2t_n r_n + 24t_n^4 u_n + 2t_n s_n u_n}{1 + t_n},
\end{aligned} \tag{3.8}$$

where $t_n = \frac{f(y_n)}{f(x_n)}$, $s_n = \frac{f(z_n)}{f(y_n)}$, $u_n = \frac{f(z_n)}{f(x_n)}$, $p_n = \frac{f(w_n)}{f(x_n)}$, $q_n = \frac{f(w_n)}{f(y_n)}$ and $r_n = \frac{f(w_n)}{f(z_n)}$.

These functions satisfy the given conditions in Theorems 1, 2, and 3, so

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - (t_n^2(5-7t_n) + (2t_n+1)(t_n^3+1) - 2t_n^4) \cdot \dfrac{f(y_n)}{f'(x_n)}, \\ w_n = z_n - ((1+s_n) + (6+u_n^2)(u_n+t_n^2) + 2(t_n-u_n)) \cdot \dfrac{f(z_n)}{f'(x_n)}, \\ x_{n+1} = w_n - \big[(1+t_n)(2t_n+t_n^2) + 3t_n^2 - t_n^3 + 8t_n^2 r_n - 4s_n^2 u_n - 2t_n^3 r_n + t_n u_n + 6t_n^2 u_n + 2(p_n+q_n) \\ \qquad + \frac{1+5u_n}{1+u_n} + \frac{2t_n^3 u_n - 10 t_n u_n^2}{1+t_n u_n} + \frac{6p_n + r_n}{1+p_n} + \frac{s_n + s_n^2 - s_n^3}{1+s_n} + \frac{2s_n u_n + 2t_n r_n + 24 t_n^4 u_n + 2 t_n s_n u_n}{1+t_n}\big] \cdot \dfrac{f(w_n)}{f'(x_n)}. \end{cases} \tag{3.9}$$

Method 7. Weight functions G, H, I, J, K, L, M, and N in (2.20) are given by

$$\begin{aligned} &G(t_n) = (1+t_n^2)(1+2t_n+2t_n^2) + t^2(2-8t_n-2t_n^2), \\ &H(t_n,s_n,u_n) = 4u_n - 5s_n + (6+s_n^3)(t_n^2+s_n) + (1+u_n^3)(1+2t_n), \\ &I(t_n) = (1+t_n)(2t_n+t_n^3) + t_n^2(4-t_n-t_n^2), \quad J(s_n) = -2s_n^2 + \frac{s_n+2s_n^2}{1+s_n^2}, \\ &K(u_n) = 1 + 6u_n - \frac{2u_n+6u_n^2}{1+u_n}, \quad L(t_n,u_n) = t_n u_n + \frac{2t_n^3 u_n - 10 t_n u_n^2 + 6 t_n^2 u_n}{1+2t_n u_n}, \\ &M(p_n,q_n,r_n) = \frac{1+2p_n+2q_n}{1-r_n} + \frac{6p_n}{1+q_n} - 1, \\ &N(t_n,s_n,u_n,r_n) = 2t_n r_n + 2s_n u_n + 24 t_n^4 u_n + \frac{6t_n^2 r_n + 6t_n^3 r_n - 4s_n^2 u_n}{1+t_n}, \end{aligned} \tag{3.10}$$

where $t_n = \frac{f(y_n)}{f(x_n)}$, $s_n = \frac{f(z_n)}{f(y_n)}$, $u_n = \frac{f(z_n)}{f(x_n)}$, $p_n = \frac{f(w_n)}{f(x_n)}$, $q_n = \frac{f(w_n)}{f(y_n)}$ and $r_n = \frac{f(w_n)}{f(z_n)}$.

These functions satisfy the given conditions in Theorems 1, 2, and 3, so

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - ((1+t_n^2)(1+2t_n+2t_n^2) + t^2(2-8t_n-2t_n^2)) \cdot \dfrac{f(y_n)}{f'(x_n)}, \\ w_n = z_n - (4u_n - 5s_n + (6+s_n^3)(t_n^2+s_n) + (1+u_n^3)(1+2t_n)) \cdot \dfrac{f(z_n)}{f'(x_n)}, \\ x_{n+1} = w_n - \big[(1+t_n)(2t_n+t_n^3) + 4t_n^2 - t_n^3 - t_n^4 - 2s_n^2 + 6u_n + 2t_n r_n + 2s_n u_n + 24 t_n^4 u_n + t_n u_n \\ \qquad + \frac{2t_n^3 u_n - 10 t_n u_n^2 + 6 t_n^2 u_n}{1+2t_n u_n} + \frac{1+2p_n+2q_n}{1-r_n} + \frac{6p_n}{1+q_n} - \frac{2u_n+6u_n^2}{1+u_n} + \frac{s_n+2s_n^2}{1+s_n^2} + \frac{6t_n^2 r_n + 6t_n^3 r_n - 4s_n^2 u_n}{1+t_n}\big] \cdot \dfrac{f(w_n)}{f'(x_n)}. \end{cases} \tag{3.11}$$

Method 8. The method by H.T. Kung and J.F. Traub [18] is given by

$$\begin{cases} y_n = x_n - \dfrac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - G_f(x_n), \\ s_n = z_n - f^2(x_n) f(y_n) H_f(x_n,y_n,z_n), \\ x_{n+1} = s_n + \frac{f^2(x_n)f(y_n)f(z_n)}{f(x_n)-f(s_n)} \left(H_f(x_n,y_n,z_n) - \frac{K_f(x_n,y_n,z_n) - L_f(x_n,y_n,z_n,s_n)}{f(x_n)-f(s_n)} \right), \end{cases} \tag{3.12}$$

where

$$
\begin{aligned}
G_f(x_n) &= \frac{f^2(x_n)f(y_n)}{f'(x_n)(f(x_n)-f(y_n))^2},\\
H_f(x_n,y_n,z_n) &= G_f(x_n)\\
&\quad \Big(\frac{-1}{f^2(x_n)(f(x_n)-f(z_n))} + \frac{f(y_n)-f(x_n)}{f(x_n)f(y_n)(f(x_n)-f(z_n))^2}\\
&\quad + \frac{1}{(f(y_n)-f(z_n))(f(x_n)-f(z_n))^2}\Big),\\
K_f(x_n,y_n,z_n) &= \frac{f(x_n)(f(y_n)-f(z_n))(f(x_n)-f(y_n)) - f^2(x_n)f(y_n)}{f'(x_n)(f(x_n)-f(z_n))(f(x_n)-f(y_n))^2(f(y_n)-f(z_n))},\\
L_f(x_n,y_n,z_n,s_n) &= \frac{G_f(x_n)(f(z_n)-f(s_n)) - (f(y_n)f^2(x_n)H_f(x_n,y_n,z_n))\,(f(y_n)-f(z_n))}{(f(y_n)-f(s_n))(f(y_n)-f(z_n))(f(z_n)-f(s_n))}.
\end{aligned}
$$

Method 9. The method by B. Neta [22] is given by

$$
\begin{cases}
y_n = x_n - \dfrac{f(x_n)}{f'(x_n)},\\
z_n = y_n - \dfrac{f(x_n)+Af(y_n)}{f(x_n)+(A-2)f(y_n)}\dfrac{f(y_n)}{f'(x_n)}, \quad A\in\mathbb{R},\\
s_n = y_n + \delta_1 f^2(x_n) + \delta_2 f^3(x_n),\\
x_{n+1} = y_n + \theta_1 f^2(x_n) + \theta_2 f^3(x_n) + \theta_3 f^4(x_n),
\end{cases}
\tag{3.13}
$$

where

$$
\begin{gathered}
F_y = f(y_n)-f(x_n), \quad F_z = f(z_n)-f(x_n), \quad F_s = f(s_n)-f(x_n),\\
\zeta_y = \frac{1}{F_y}\left(\frac{y_n-x_n}{F_y} - \frac{1}{f'(x_n)}\right), \zeta_z = \frac{1}{F_z}\left(\frac{z_n-x_n}{F_z} - \frac{1}{f'(x_n)}\right), \zeta_s = \frac{1}{F_s}\left(\frac{s_n-x_n}{F_s} - \frac{1}{f'(x_n)}\right),\\
\delta_1 = \zeta_y + \delta_2 F_y, \qquad \delta_2 = -\frac{\zeta_y-\zeta_z}{F_y-F_z}, \qquad \gamma_1 = \frac{\zeta_s-\zeta_z}{F_s-F_z}, \qquad \gamma_2 = \frac{\zeta_y-\zeta_z}{F_y-F_z},\\
\theta_1 = \zeta_s + \theta_2 F_s - \theta_3 F_s^2, \quad \theta_2 = \gamma_1 + \theta_3(F_s - +F_z), \quad \theta_3 = \frac{\gamma_1-\gamma_2}{F_s-F_y},
\end{gathered}
$$

Method 10. The method by Y. H. Geum and Y. I. Kim [11] is given by

$$
\begin{cases}
y_n = x_n - \dfrac{f(x_n)}{f'(x_n)},\\
z_n = y_n - K_f(u_n)\cdot\dfrac{f(y_n)}{f'(x_n)},\\
s_n = z_n - H_f(u_n,v_n,w_n)\cdot\dfrac{f(z_n)}{f'(x_n)},\\
x_{n+1} = s_n - W_f(u_n,v_n,w_n,t_n)\cdot\dfrac{f(s_n)}{f'(x_n)}.
\end{cases}
\tag{3.14}
$$

where

$$K_f(u_n) = \frac{1+\beta u_n+\lambda u_n^2}{1+(\beta-2)u_n+\mu u_n^2},$$
$$H_f(u_n,v_n,w_n) = \frac{1+au_n+bv_n+\gamma w_n}{1+cu_n+dv_n+\sigma w_n},$$
$$W_f(u_n,v_n,w_n,t_n) = \frac{1+B_1u_n+B_2v_nw_n}{1+B_3v_n+B_4w_n+B_5t_n+B_6v_nw_n}+G(u_n,w_n),$$

and $u_n=\frac{f(y_n)}{f(x_n)}$, $v_n=\frac{f(z_n)}{f(y_n)}$, $w_n=\frac{f(z_n)}{f(x_n)}$ and $t_n=\frac{f(s_n)}{f(z_n)}$.

$$a=2,\quad b=0,\quad c=0,\quad d=-1,\quad \gamma=2+\sigma,\quad \lambda=-9+\frac{5\beta}{2},\quad \mu=-4+\frac{\beta}{2},$$
$$B_1=2,\quad B_2=2+\sigma,\quad B_3=-1,\quad B_4=-2,\quad B_5=-1,\quad B_6=2(1+\sigma),$$

With weight function

$$\begin{aligned} G(u_n,w_n) &= \frac{-1}{2}\left(u_nw_n\left(6+12u_n+u_n^2(24-11\beta)+u_n^3\phi_1+4\sigma\right)\right)+\phi_2w_n^2,\\ \beta&=2,\quad \sigma=-2,\quad \phi_1=11\beta^2-66\beta+136,\quad \phi_2=2u_n(\sigma^2-2\sigma-9)-4\sigma-6. \end{aligned} \tag{3.15}$$

Method 11. The method by Y. H. Geum and Y. I. Kim [12] is given by

$$\begin{cases} y_n = x_n-\dfrac{f(x_n)}{f'(x_n)},\\ z_n = y_n-K_f(u_n)\cdot\dfrac{f(y_n)}{f'(x_n)},\\ s_n = z_n-H_f(u_n,v_n,w_n)\cdot\dfrac{f(z_n)}{f'(x_n)},\\ x_{n+1} = s_n-W_f(u_n,v_n,w_n,t_n)\cdot\dfrac{f(s_n)}{f'(x_n)}. \end{cases} \tag{3.16}$$

where

$$K_f(u_n) = \frac{1+\beta u_n+\gamma u_n^2}{1+(\beta-2)u_n+\mu u_n^2},$$
$$H_f(u_n,v_n,w_n) = \frac{1+au_n+bv_n+\gamma w_n}{1+cu_n+dv_n+\sigma w_n},$$
$$W_f(u_n,v_n,w_n,t_n) = \frac{1+A_1u_n}{1+A_2v_n+A_3w_n+A_4t_n}+G(u_n,v_n,w_n),$$

also $u_n=\frac{f(y_n)}{f(x_n)}$, $v_n=\frac{f(z_n)}{f(y_n)}$, $w_n=\frac{f(z_n)}{f(x_n)}$ and $t_n=\frac{f(s_n)}{f(z_n)}$,

$$a=2,\quad b=0,\quad c=0,\quad d=-1,$$
$$\gamma=2+\sigma,\quad \lambda=-9+\frac{5\beta}{2},\quad \mu=-4+\frac{\beta}{2},$$
$$A_1=2,\quad A_2=-1,\quad A_3=-2,\quad A_4=-1,$$

with the weight function

$$\begin{aligned} G(u_n,v_n,w_n) &= -6u_n^3v_n+6w_n^2-4u_n^4(3v_n+17w_n)+u_n(2v_n^2+4v_n^3+w_n-2w_n^2),\\ \beta&=0,\quad \sigma=-2. \end{aligned} \tag{3.17}$$

We test our proposed methods (3.7), (3.9), and (3.11) on the functions $f_1, \ldots, f_6$ described in Table 1, the Table also lists the exact roots x^* and initial approximations x_0, which are computed using the `FindRoot` command of Mathematica [14, pp. 158–160].

For the methods defined by (3.7), (3.9), and (3.11), we display in Tables 2-4 for the weight functions (3.6), (3.8), (3.10) the errors $|x_n - x^*|$ for $n = 1, 2, 3$, and the *computational order of convergence (coc)* [35], approximated by

$$\text{coc} \approx \frac{\ln |(x_{n+1} - x^*)/(x_n - x^*)|}{\ln |(x_n - x^*)/(x_{n-1} - x^*)|}.$$

Test function f_n	root x^*	initial approximation x_0
$f_1(x) = \ln(1 + x^2) + e^x \sin x$	0	0.03
$f_2(x) = \frac{-x}{100} + \sin x$	0	0.5
$f_3(x) = x \ln(1 + x \sin x) + e^{-1+x^2+x\cos x} \sin \pi x$	0	0.01
$f_4(x) = 1 + e^{2+x-x^2} + x^3 - \cos(1 + x)$	-1	-0.3
$f_5(x) = (1 - \sin x^2)\frac{x^2+1}{x^3+1} + x \ln(x^2 - \pi + 1) - \frac{1+\pi}{1+\sqrt{\pi^3}}$	$\sqrt{\pi}$	1.7
$f_6(x) = (1 + x^2)\cos(\frac{\pi x}{2}) + \frac{\ln(x^2+2x+2)}{1+x^2}$	-1	-1.1

Table 1: Test functions $f_1, \ldots, f_6$, root x^* and initial approximation x_0.

To obtain a high accuracy and avoid loss of significant digits, we employ multi-precision arithmetic with 6000 significant decimal digits in Mathematica 8.

f_n	$\lvert x_1 - x^*\rvert$	$\lvert x_2 - x^*\rvert$	$\lvert x_3 - x^*\rvert$	coc
f_1	$0.380e-20$	$0.126e-319$	$0.276e-5111$	16.0000
f_2	$0.104e-10$	$0.265e-192$	$0.211e-3279$	17.0000
f_3	$0.450e-28$	$0.303e-449$	$0.561e-7188$	16.0000
f_4	$0.609e-8$	$0.465e-136$	$0.630e-2186$	16.0000
f_5	$0.246e-14$	$0.276e-230$	$0.169e-3685$	16.0000
f_6	$0.142e-17$	$0.482e-283$	$0.139e-4530$	16.0000

Table 2: Errors and coc for method (3.7).

f_n	$\lvert x_1 - x^*\rvert$	$\lvert x_2 - x^*\rvert$	$\lvert x_3 - x^*\rvert$	coc
f_1	$0.144e-19$	$0.193e-310$	$0.222e-4964$	16.0000
f_2	$0.301e-23$	$0.339e-451$	$0.336e-8582$	19.0000
f_3	$0.405e-28$	$0.515e-450$	$0.239e-7200$	16.0000
f_4	$0.628e-8$	$0.276e-135$	$0.561e-2173$	16.0000
f_5	$0.224e-14$	$0.526e-231$	$0.456e-3697$	16.0000
f_6	$0.186e-17$	$0.322e-281$	$0.202e-4501$	16.0000

Table 3: Errors and coc for method (3.9).

Tables 2-4 show that methods (3.7), (3.9), and (3.11) support the convergence analysis given in the previous sections.

f_n	$\lvert x_1 - x^* \rvert$	$\lvert x_2 - x^* \rvert$	$\lvert x_3 - x^* \rvert$	coc
f_1	$0.389e-20$	$0.931e-321$	$0.107e-5130$	16.0000
f_2	$0.414e-22$	$0.1220e-385$	$0.117e-6565$	17.0000
f_3	$0.936e-29$	$0.865e-461$	$0.243e-7373$	16.0000
f_4	$0.554e-8$	$0.756e-136$	$0.108e-2181$	16.0000
f_5	$0.119e-14$	$0.116e-235$	$0.760e-3772$	16.0000
f_6	$0.125e-17$	$0.926e-285$	$0.745e-4559$	16.0000

Table 4: Errors and coc for method (3.11) .

In Tables 5-7, we compare our three-point method (3.2) with the methods (3.3), (3.4) and (3.5) and our four-point methods (3.7), (3.9) and (3.11) with the methods (3.12), (3.13), (3.14) and (3.16).

Method	Weight function	$\lvert x_1 - x^* \rvert$	$\lvert x_2 - x^* \rvert$	$\lvert x_3 - x^* \rvert$	coc
(3.2)	(3.1)	$0.834e-13$	$0.112e-121$	$0.160e-1101$	9.0000
(3.3)	-	$0.316e-11$	$0.165e-95$	$0.918e-770$	8.0000
(3.4)	-	$0.445e-11$	$0.380e-94$	$0.108e-758$	8.0000
(3.5)	-	$0.704e-11$	$0.658e-92$	$0.384e-740$	8.0000
(3.7)	(3.6)	$0.844e-25$	$0.659e-435$	$0.257e-7406$	17.0000
(3.9)	(3.8)	$0.102e-24$	$0.229e-433$	$0.215e-7380$	17.0000
(3.11)	(3.10)	$0.371e-25$	$0.429e-465$	$0.571e-8384$	18.0000
(3.12)	-	$0.114e-22$	$0.323e-374$	$0.546e-5999$	16.0000
(3.13)	-	$0.226e-22$	$0.403e-369$	$0.429e-5917$	16.0000
(3.14)	(3.15)	$0.235e-22$	$0.106e-368$	$0.316e-5910$	16.0000
(3.16)	(3.17)	$0.310e-21$	$0.600e-350$	$0.226e-5609$	16.0000

Table 5: Comparison for $f(x) = \ln(1 - x + x^2) + 4\sin(1 - x)$, zero $x^* = 1$ and initial $x_0 = 1.1$.

Method	Weight function	$\lvert x_1 - x^* \rvert$	$\lvert x_2 - x^* \rvert$	$\lvert x_3 - x^* \rvert$	coc
(3.2)	(3.1)	$0.125e-10$	$0.888e-85$	$0.545e-678$	8.0000
(3.3)	-	$0.414e-9$	$0.961e-72$	$0.811e-573$	8.0000
(3.4)	-	$0.597e-9$	$0.273e-70$	$0.533e-561$	8.0000
(3.5)	-	$0.629e-9$	$0.393e-70$	$0.929e-560$	8.0000
(3.7)	(3.6)	$0.201e-9$	$0.510e-148$	$0.142e-2365$	16.0000
(3.9)	(3.8)	$0.210e-9$	$0.807e-148$	$0.183e-2362$	16.0000
(3.11)	(3.10)	$0.389e-20$	$0.931e-321$	$0.107e-5130$	16.0000
(3.12)	-	$0.883e-18$	$0.897e-282$	$0.114e-4505$	16.0000
(3.13)	-	$0.183e-17$	$0.254e-276$	$0.470e-4418$	16.0000
(3.14)	(3.15)	$0.274e-10$	$0.682e-162$	$0.150e-2587$	16.0000
(3.16)	(3.17)	$0.115e-9$	$0.377e-149$	$0.621e-2381$	16.0000

Table 6: Comparison for $f(x) = \ln(1 + x^2) + e^x \sin x$, zero $x^* = 0$ and initial $x_0 = 0.1$.

Method	Weight function	$\lvert x_1 - x^* \rvert$	$\lvert x_2 - x^* \rvert$	$\lvert x_3 - x^* \rvert$	coc
(3.2)	(3.1)	$0.361e-13$	$0.209e-106$	$0.270e-852$	8.0000
(3.3)	-	$0.230e-12$	$0.396e-99$	$0.303e-793$	8.0000
(3.4)	-	$0.349e-12$	$0.168e-97$	$0.485e-780$	8.0000
(3.5)	-	$0.477e-17$	$0.463e-141$	$0.366e-1133$	8.0000
(3.7)	(3.6)	$0.499e-25$	$0.113e-400$	$0.536e-6411$	16.0000
(3.9)	(3.8)	$0.406e-25$	$0.345e-402$	$0.256e-6435$	16.0000
(3.11)	(3.10)	$0.825e-26$	$0.242e-414$	$0.760e-6631$	16.0000
(3.12)	-	$0.211e-24$	$0.152e-390$	$0.780e-6249$	16.0000
(3.13)	-	$0.484e-24$	$0.209e-384$	$0.310e-6150$	16.0000
(3.14)	(3.15)	$0.253e-24$	$0.173e-389$	$0.372e-6232$	16.0000
(3.16)	(3.17)	$0.454e-22$	$0.108e-350$	$0.123e-5608$	16.0000

Table 7: Comparison for $f(x) = \frac{-2}{27}(9\sqrt{2} + 7\sqrt{3}) + \sqrt{1-x^2} + (1+x^3)\cos(\frac{\pi x}{2})$, zero $x^* = \frac{1}{3}$ and initial $x_0 = 0.35$.

It can be observed from Tables 5-7 that for the presented examples our three-point proposed method (3.2) is comparable and competitive to the methods (3.3), (3.4) and (3.5) and our four-point proposed methods (3.7), (3.9) and (3.11) are comparable and competitive to the methods (3.12), (3.13), (3.14) and (3.16) also.

3.2 Dynamic behavior

In this section, we survey the comparison of iterative methods in the complex plane by using basins of attraction. Studying the dynamic behavior, using basins of attractions, of the rational functions associated to an iterative method gives important information about the convergence and stability of the scheme [31].

Neta et al. have compared various methods for solving nonlinear equations with multiple roots by comparing the basins of attraction [24] and also Scott et al. have compared several methods for approximating simple roots [29]. Moreover, a number of iterative root-finding methods were compared from a dynamical point of view by Amat et al. [1]-[4], Neta et al. [23], [25] Stewart [32], Vrscay and Gilbert [34]. To this end, some basic concepts are briefly recalled.

Let $G : \mathbb{C} \to \mathbb{C}$ be a rational map on the complex plane. For $z \in \mathbb{C}$, we define its orbit as the set $orb(z) = \{z, G(z), G^2(z), \dots\}$. A point $z_0 \in \mathbb{C}$ is called a periodic point with minimal period m if $G^m(z_0) = z_0$, where m is the smallest integer with this property. A periodic point with minimal period 1 is called a fixed point. Moreover, a point z_0 is called attracting if $|G'(z_0)| < 1$, repelling if $|G'(z_0)| > 1$, and neutral otherwise. The Julia set of a nonlinear map $G(z)$, denoted by $J(G)$, is the closure of the set of its repelling periodic points. The complement of $J(G)$ is the Fatou set $F(G)$, where the basin of attraction of the different roots lie [5], [7].

We use the basin of attraction for comparing the iteration algorithms. Approximating basins of attraction is a method to visually comprehend how an algorithm behaves as a function of the various starting points.

From the dynamical point of view, in fact, we take a 256×256 grid of the square $[-3, 3] \times [-3, 3] \in \mathbb{C}$ and assign a color to each grid point z_0 according to the simple root to which the corresponding orbit of the iterative method starting from z_0 converges, and we mark the point as black if the orbit does not converge to a root, in the sense that after at most 100 iterations it has a distance to any of the roots, which is larger than 10^{-3}. In this way, we distinguish the attraction basins by their color for different methods.

Test Problems p_n	Roots		
$p_1(z) = z^2 + 1$	i,	$-i$	
$p_2(z) = z^3 + z$	0,	i,	$-i$
$p_3(z) = z^3 + z^2 - 1$	$-0.877439 + 0.744862i$,	$-0.877439 - 0.744862i$,	0.7548878

Table 8: Test Problems $p_1(z), p_2(z), p_3(z)$ and roots.

For the test problem $p_1(z)$, with its roots given in Table 8, the results are presented in Figures 1-5. For test problems $p_2(z)$ and $p_3(z)$, the results are shown in Figures 6-10 and Figures 11-15, respectively. As a result, the method (3.9) (see Figures 1, 6 and 11) seems to produce larger basins of attraction than the methods (3.14) and (3.16) (see Figures 4, 5, 9, 10, 14 and 15) and smaller basins of attraction than the methods (3.12) and (3.13) (see Figures 2, 3, 7, 8, 12 and 13).

Note that points might belong to no basin of attraction; these are starting points for which the methods do not converge, approximated and visualized by black points. These exceptional points constitute the so-called Julia set of the corresponding methods.



Figure 1: Method (3.9) for test problem p_1



Figure 2: Method (3.12) for test problem p_1



Figure 3: Method (3.13) for test problem p_1



Figure 4: Method (3.14) for test problem p_1



Figure 5: Method (3.16) for test problem p_1



Figure 6: Method (3.9) for test problem p_2



Figure 7: Method (3.12) for test problem p_2



Figure 8: Method (3.13) for test problem p_2



Figure 9: Method (3.14) for test problem p_2



Figure 10: Method (3.16) for test problem p_2



Figure 11: Method (3.9) for test problem p_3



Figure 12: Method (3.12) for test problem p_3



Figure 13: Method (3.13) for test problem p_3



Figure 14: Method (3.14) for test problem p_3



Figure 15: Method (3.16) for test problem p_3

4 Conclusion

We introduce a new optimal class of four-point methods without memory for approximating a simple root of a given nonlinear equation. Our proposed methods use five function evaluations for each iteration. Therefore they support Kung and Traub's conjecture. Error and convergence analysis are carried out. Numerical examples show that our methods work and can compete with other methods in the same class. We used the basin of attraction for comparing the iteration algorithms.

References

[1] Amat, S., Busquier, S., Plaza, S., Iterative root-finding methods, Unpublished report, (2004).

[2] Amat, S., Busquier, S., Plaza, S., Review of some iterative root-finding methods from a dynamical point of view, Scientia Series A: Math. Sci., 10, 3-35, (2004).

[3] Amat, S., Busquirer, S., Plaza, S., Dynamics of a family of third-order itrative methods that do not require using second derivatives, Appl. Math. Comput., 154, 735-746, (2004).

[4] Amat, S., Busquier, S., Plaza, S., Dynamics of the King and Jarratt iterations, Aeq. Math., 69, 212-223, (2005).

[5] Babajee, D.K.R., Thukral, R., On a 4-point sixteen-oredr King family of iterative method for solving nonlinear equations, International Journal of Mathematics and Mathematical Sciences, DOI: 10.1155/2012/979245, (2012).

[6] Chun, C., Lee, M.Y., A new optimal eighth-order family of iterative methods for the solution of nonlinear equations, Appl. Math. Comput., 223, 506-519, (2013).

[7] Cordero, A., Fardi, M., Ghasemi, M., Torregrosa, J.R., Accelerated iterative methods for finding solutions of nonlinear equations and their dynamical behavior, Calcolo, 1-14, (2012).

[8] Cordero, A., Lotfi, T., Bakhtiari, P., Torregrosa, J.R., An efficient two-parametric family with memory for nonlinear equations, Numer. Algor., DOI 10.1007/s11075-014-9846-8, (2014).

[9] Cordero, A., Lotfi, T., Mahdiani, K., Torregrosa, J.R., Two optimal general classes of iterative methods with eighth-order, Acta Appl. Math., DOI 10.1007/s10440-014-9869-0, (2013).

[10] Cordero, A., Torregrosa, J.R., Vassileva, M.P., Three-step iterative methods with optimal eighth-order convergence, J. Comput. Appl. Math., 235, 3189-3194, (2011).

[11] Geum, Y. H., Kim, Y. I., A biparametric family optimally convergent sixteenth-order multipoint methods with their fourth-step weighting function as a sum of a rational and a generic two-variable function, J. Comput. Appl. Math., 235, 3178-3188, (2011).

[12] Geum, Y. H., Kim, Y. I., A family of optimal sixteen-order multipoint method with a linear fraction plus a trivariate polynomial as the fourth-step weighting function, Computers, Math. with Appl., 61, 3278-3287, (2011).

[13] Geum, Y. H., Kim, Y. I., A biparametric family of four-step sixteen-oredr root-finding methods with the optimal efficiency index, Appl. Math. Lett., 24, 1336-1342, (2011).

[14] Hazrat, R., Mathematica: A Problem-Centered Approach, Springer-Verlag, 2010.

[15] Jarratt, P., Some fourth order multipoint iterative methods for solving equations, Math. Comput., 20, 434-437, (1966).

[16] Khattri, S.K., Steihaug, T., Algorithm for forming derivative-free optimal methods, Numer. Algor., 65, 809-824, (2014).

[17] King, R.F., Family of four order methods for nonlinear equations SIAM J. Numer. Anal., 10, 876-879, (1973).

[18] Kung, H.T., Traub, J.F., Optimal order of one-point and multipoint iteration, Assoc. Comput. Math. 21, 643-651, (1974).

[19] Lotfi, T., Cordero, A., Torregrosa, J.R., Amir Abadi, M., Mohammadi Zadeh, M., On generalization based on Bi et al. iterative methods with eighth-order convergence for solving nonlinear equations, The Scientific World Journal, Article ID 272949, (2014).

[20] Lotfi, T., Sharifi, S., Salimi, M., Siegmund, S., A new class of three-point methods with optimal convergence order eight and its dynamics, Numer. Algor., DOI 10.1007/s11075-014-9843-y, (2014).

[21] Lotfi, T., Tavakoli, E., On a new efficient Steffensen-like iterative class by applying a suitable self-accelerator parameter, The Scientific World Journal, Article ID 769758, (2014).

[22] Neta, B., On a family of multipoint methods for nonlinear equations, Intern. J. Computer Math., 9, 353-361, (1981).

[23] Neta, B., Chun, C., Scott, M., Basin of attractions for optimal eighth order methods to find simpl roots of nonlinear equations, Appl. Math. Comput., 227, 567-592, (2014).

[24] Neta, B., Scott, M., Chun, C., Basin attractors for various methods for multiple roots, Appl. Math. Comput., 218, 5043-5066, (2012).

[25] Neta, B., Scott, M., Chun, C., Basin of attraction for several methods to find simple roots of nonlinear equations, Appl. Math. Comput., 218, 10548-10556, (2012).

[26] Ostrowski, A.M., Solution of Equations and Systems of Equations, Academic Press, New York, 1966.

[27] Petkovic, M.S., A note on the priority of optimal multipoint methods for solving nonlinear equations, Appl. Math. Comput., 219, 5249-5252, (2013).

[28] Petkovic, M.S., Neta, B., Petkovic, L.D., Dzunic, J., Multipoint methods for solving nonlinear equations: A survey, Appl. Math. Comput., 226, 635-660, (2014).

[29] Scott, M., Neta, B., Chun, C., Basin attractors for various methods, Appl. Math. Comput., 218, 2584-2599, (2011).

[30] Sharma, J.R., Sharma, R., A new family of modified Ostrowski's methods with accelerated eighth order convergence, Numer. Algor., 54, 445-458, (2010).

[31] Soleymani, F., Lotfi, T., Bakhtiari, P., A multi-step class of iterative methods for nonlinear systems, Optim. Lett., 8(3), 1001-1015, (2014).

[32] Stewart, B.D., Attractor Basins of Various Root-Finding Methods M.S. thesis, Naval Postgraduate School, Department of Applied Mathematics, Monterey, CA, June, 2001.

[33] Traub, J.F., Iterative Methods for the Solution of Equations, Prentice Hall, New York, 1964.

[34] Vrscay, E.R., Gilbert, W.J., Extraneous fixed points, basin boundaries and chaotic dynamics for Schroder and Konig rational iteration functions, Numer. Math., 52, 1-16, (1988).

[35] Weerakoon, S., Fernando, T.G.I., A variant of Newton's method with accelerated third-order convergence, Appl. Math. Lett., 13, 87-93, (2000).